

# Status assessment of raptors in Cape Verde confirms a major crisis for scavengers

SABINE M. HILLE and NIGEL J. COLLAR

**Abstract** Scavenging raptors have been postulated to be declining at a rate far higher than predatory raptors. To test this hypothesis we reviewed the historical and present status of the seven raptor species—three scavengers (two kites and a vulture), one partial scavenger (a buzzard) and three species (osprey and two falcons) that take live prey—that breed on the Cape Verde islands. Scavenging raptors have experienced steeper declines and more local extinctions than non-scavengers in Cape Verde, with the partial scavenger midway between the two groups. Causes of scavenger decline include incidental poisoning, direct persecution and declines in the availability of carcasses and other detritus. These findings, which highlight the conservation importance of the island of Santo Antão, indicate the priority that needs to be accorded to scavengers, particularly in Europe where many insular populations are reaching unsustainable levels.

**Keywords** Cape Verde, decline rates, poisoning, persecution, raptor community, vultures

## Introduction

The decline of populations of scavenging raptors, notably vultures and kites, is emerging as a serious phenomenon. Most dramatic has been the loss of > 90% of *Gyps* vultures in the Indian subcontinent as a result of accidental diclofenac poisoning (Pain et al., 2008). However, this extraordinary circumstance has overshadowed events elsewhere, although in some respects these are more worrying because they are less obviously remediable. There have been steep declines in South-East Asia's vulture populations (Pain et al., 2008; J. W. Duckworth et al., unpubl. data). West Africa has lost 98% of its large vultures outside protected areas within 30 years (Thiollay, 2006). Greece lost 84% of its bearded vultures *Gypaetus barbatus* within 10 years (Xirouchakis & Tsiakiris, 2009). The Egyptian *Neophron percnopterus* and red-headed vulture *Sarcogyps calvus* were categorized as Endangered and Critically Endangered, respectively, on the IUCN Red List

in 2007, following assembly of the evidence (Cuthbert et al., 2006). Across southern Europe the red kite *Milvus milvus* is declining in all countries where it occurs (BirdLife International, 2009). The black kite *Milvus migrans*, although generally widespread and common, is also showing signs of decline (Sergio et al., 2003). The deteriorating status of these species seems primarily to be linked to incidental mortality from feeding on carcasses deliberately poisoned to control mammals regarded as pests, although direct persecution, disturbance to breeding birds and diminution of food supplies are also believed to play a part (see, for example, accounts of scavengers in Tucker & Heath, 1994).

Raptors in Cape Verde comprise seven species: black kite, red kite, Egyptian vulture and common buzzard *Buteo buteo* (Accipitridae), osprey *Pandion haliaetus* (Pandionidae), and common kestrel *Falco tinnunculus* and peregrine *Falco peregrinus* (Falconidae). Four of these species occur as five endemic subspecies, *fasciicauda* (red kite), *bannermani* (common buzzard), *neglectus* and *alexandri* (common kestrel) and *madens* (peregrine), all of which Hazevoet (1995) elevated to species under the phylogenetic species concept but which we here retain as subspecies. Three of these species, the two kites and vulture, may be typified as scavengers, as this is their primary mode of foraging (all take live prey opportunistically), although the buzzard often also scavenges; the others exclusively take live prey. All seven are represented strongly in Europe, and the Cape Verde populations of osprey, red kite, common buzzard and common kestrel are the south-westernmost outliers of their Palaearctic ranges, and there are Afrotropical breeding populations of black kite, Egyptian vulture and peregrine.

The vulnerability of scavengers has been detected in a general analysis of avian susceptibility to environmental change, with a prediction of relatively high losses of members of this guild in the 21st century (Sekercioglu et al., 2004). Using the Cape Verde raptor assemblage as a case study we examine whether the patterns of population change in the archipelago's seven species confirm other evidence that scavengers are faring worse than birds that catch live prey.

## Methods

For each species we synthesized relevant information (status assessments and individual records) from all available literature, published and unpublished, and arranged it by species and island (13 in the Cape Verde archipelago,

SABINE M. HILLE (Corresponding author) Institute for Wildlife Biology and Game Management, University of Natural Resources and Life Sciences, Vienna, Austria. E-mail [sabine.hille@boku.ac.at](mailto:sabine.hille@boku.ac.at)

NIGEL J. COLLAR BirdLife International, Girton Road, Cambridge, UK

Received 8 January 2010. Revision requested 17 May 2010.  
Accepted 11 June 2010.

with Ilhéus do Rombo treated as one island: Fig. 1). We added unpublished information from fieldwork by SMH in 1996–2003 involving (1) a 1997 census of osprey and Egyptian vulture on Santo Antão, São Vicente, Boavista, Sal, Brava, Fogo and Maio; (2) a 2001 census and territory mapping of all raptors across Santo Antão in which carrion feeders were lured with goat carcasses; (3) a 2001 census of all raptors on Santa Luzia, Raso and Branco (with Raso censused again in 2003); (4) a 2002 month-long survey of kites on Maio and Boavista; (5) an intensive search for all nest sites of common kestrels in 1996–2001 in which 30% of every island's surface, representing the main environmental units (desert, agricultural field and settlements), were covered (only breeding pairs counted), with extrapolation of total nests based on proportions of these units per island. Because habitat quality varies strongly both between and within islands we did not calculate population densities for each species per island.

We then filtered this material for highest abundance (often qualitative, especially in accounts by earlier visitors) and most recent abundance. We used these two values to tabulate change over time and classified populations crudely into three status conditions: common (taxa that, according to statements in the literature or to observations by SMH, can or could regularly be seen in most parts of a given island), rare (taxa not numerous and restricted to certain areas of an island), and extinct, which includes vagrants (the legend to Table 1 provides further definitions of these categories). We then assessed status trends by depicting changes between these three status conditions by island and species: no change was classed as such, a one-category change (e.g. common to rare) was classed as a decline, and a two-category change (e.g. common to extinct) was classed as severe decline. Each island was assigned one of these trend categories to show the proportion of trend categories per species.




FIG. 1 The Cape Verde archipelago. The inset shows the location of the archipelago off the west coast of Africa.

TABLE 1 Distribution and population trends of Cape Verde raptors by island, setting highest reported abundance against most recent evidence. Each species has an upper (highest numerical value) and lower (latest numerical value) row. Under each island a number or code (abd, abundant; br, breeding; com, common; n/r, no record; num, numerous; p, pairs; pres, present; unk, unknown; wds, widespread; ?, situation uncertain), with superscript reference or comment, is followed by a status letter (C, common: generally a number of descriptions suggesting island well stocked, interpreted as easy to find; R, rare: generally < 5 individuals, interpreted as hard to find; X, extinct; X?, situation uncertain: rarely present, vagrant, not proven to breed, likely to be inviable or probably extinct). For common kestrel: *n*, subspecies *neglectus*; *a*, subspecies *alexandri*.

Species	Values	Santo Antão	São Vicente	Santa Luzia	Branco	Raso	São Nicolau	Sal	Boavista	Maio	Santiago	Fogo	Ilhéus do Rombo	Brava
Black kite <i>Milvus migrans</i>	Highest number + status Latest record + status	num <sup>1</sup> C 1 <sup>9</sup> X?	3–5 <sup>2</sup> R 1 <sup>10</sup> X?	br <sup>3</sup> R 0 <sup>11</sup> X	1 <sup>4</sup> X? unk X?	br <sup>3</sup> R 0 <sup>12</sup> X?	com <sup>5,6</sup> C 0 <sup>12</sup> X?	br <sup>7</sup> R 0 <sup>12</sup> X?	br <sup>3</sup> R 1 <sup>13</sup> X?	abd <sup>3</sup> R 11 <sup>12</sup> X?	abd <sup>3</sup> C 1 <sup>12</sup> X?	br <sup>3</sup> C 0 <sup>11</sup> X	? <sup>8,3</sup> X? ? <sup>11,14,15</sup> X?	br <sup>3</sup> R 0 <sup>12</sup> X
Red kite <i>Milvus milvus</i>	Highest number + status Latest record + status	5–7 <sup>12,16</sup> R 0 <sup>12</sup> X	com <sup>17</sup> R 2 <sup>16</sup> X	n/r n/r	– <sup>4</sup> R – <sup>4</sup> X	– <sup>4</sup> R – <sup>4</sup> X	>12 <sup>11</sup> C 0 <sup>12,21,22</sup> X	n/r n/r	n/r <sup>18</sup> n/r <sup>18</sup>	n/r <sup>18</sup> n/r <sup>18</sup>	num <sup>19</sup> R 0 <sup>12,21,22</sup> X	n/r n/r	1 <sup>11,20</sup> X? 0 X?	com <sup>17</sup> R 0 <sup>11,12</sup> X
Egyptian vulture <i>Neophron percnopterus</i>	Highest number + status Latest record + status	com <sup>15</sup> C 10 p <sup>12</sup> C <sup>25</sup>	50+ <sup>17</sup> C 0 <sup>12</sup> X?	1–2 <sup>8</sup> R 0 <sup>12</sup> X	pres? <sup>17</sup> X? unk X?	br <sup>11</sup> R 0 <sup>12</sup> X	abd <sup>23</sup> C 0 <sup>12</sup> R <sup>25</sup>	br <sup>3</sup> R 0 <sup>12</sup> X	br <sup>8,24</sup> R 2–3 p <sup>12</sup> R	2 <sup>15</sup> R 0 <sup>12</sup> X?	wds <sup>8,15,17</sup> C 0 <sup>12</sup> X	1–2 p <sup>17</sup> C 0 <sup>12</sup> X?	1–2 p <sup>3</sup> R unk X?	wds <sup>8</sup> R 0 <sup>12</sup> X?
Common buzzard <i>Buteo buteo</i>	Highest number + status Latest record + status	11–13 p <sup>12</sup> R 11–13 p <sup>12</sup> R	1 <sup>15,17</sup> R 0 <sup>11</sup> X	n/r n/r	n/r n/r	n/r n/r	5 <sup>26</sup> R 1 <sup>16,28</sup> X?	n/r n/r	pres <sup>8</sup> R 1 <sup>16,29</sup> X	n/r n/r	20–30 p <sup>27</sup> R 3 <sup>30</sup> R <sup>25</sup>	1 <sup>3</sup> R 0 <sup>12</sup> X	n/r n/r	1 <sup>1</sup> R 0 <sup>12</sup> X
Osprey <i>Pandion haliaetus</i>	Highest number + status Latest record + status	18–23 p <sup>31</sup> C 18–23 p <sup>31</sup> C	7–9 p <sup>32</sup> C 8 p <sup>31</sup> C	5–6 p <sup>31</sup> C 5–6 p <sup>31</sup> C	3–4 p <sup>33</sup> C 1–2 p <sup>12,31</sup> R <sup>25</sup>	5–6 p <sup>12,30</sup> C <sup>34</sup> 5–6 p <sup>12,30</sup> C <sup>34</sup>	17 p <sup>31</sup> C 17 p <sup>31</sup> C	6–8 p <sup>33,35</sup> C 4 p <sup>31</sup> R <sup>37</sup>	17 ± 2 p <sup>36</sup> C 17 ± 2 p <sup>36</sup> C <sup>34</sup>	2–3 p <sup>3</sup> R 1 p <sup>31</sup> R <sup>25</sup>	4–6 p <sup>33</sup> R 3–4 p <sup>31</sup> R	2–5 p <sup>3</sup> C 0 <sup>33</sup> X	br <sup>3,33</sup> R 0 <sup>12,31</sup> X	num <sup>23</sup> C 0 <sup>31</sup> X
Common kestrel <i>Falco tinnunculus</i>	Highest number + status Latest record + status	<i>n</i> 400–500 p <sup>12</sup> C <i>n</i> 400–500 p <sup>12</sup> C	<i>n</i> 50 p <sup>12</sup> R <i>n</i> 50 p <sup>12</sup> R	<i>n</i> 4 p <sup>12</sup> C <i>n</i> 4 p <sup>12</sup> C	<i>n</i> 1–2 p <sup>11</sup> R <i>n</i> 1 p <sup>12</sup> R	<i>n</i> 1–2 p <sup>11</sup> R <i>n</i> 1 p <sup>12</sup> R	<i>n</i> 250 p <sup>12</sup> C <i>n</i> 250 p <sup>12</sup> C	<i>a</i> 40 p <sup>12</sup> C <i>a</i> 40 p <sup>12</sup> C	<i>a</i> 150 p <sup>12,38</sup> C <i>a</i> 150 p <sup>12,38</sup> C	<i>a</i> 50 p <sup>12</sup> C <i>a</i> 50 p <sup>12</sup> C	<i>a</i> 400 p <sup>12</sup> C <i>a</i> 400 p <sup>12</sup> C	<i>a</i> 250 p <sup>12</sup> C <i>a</i> 250 p <sup>12</sup> C	<i>a</i> 1–2 p <sup>11</sup> R <i>a</i> 1 p <sup>12</sup> R	<i>a</i> 60 p <sup>12</sup> C <i>a</i> 60 p <sup>12</sup> C
Peregrine <i>Falco peregrinus</i>	Highest number + status Latest record + status	2 p <sup>12</sup> R 2 p <sup>12</sup> R	1 <sup>11</sup> R 1 <sup>11</sup> R	n/r n/r	1 <sup>11</sup> R 0 <sup>40</sup> X?	pres <sup>16</sup> R pres <sup>16</sup> R	1 <sup>11</sup> R pres <sup>16</sup> R	2 <sup>2</sup> R 2 <sup>2</sup> R	pres <sup>12,16</sup> R pres <sup>12,16</sup> R	n/r n/r	4 p <sup>26</sup> R 2+ p <sup>30</sup> R	1 <sup>26</sup> R pres <sup>16</sup> R	br <sup>3</sup> X? pres <sup>26</sup> X?	br <sup>39</sup> R 1 <sup>12</sup> R

<sup>1</sup>Nørrevang & den Hartog (1984). <sup>2</sup>Hazevoet (1998). <sup>3</sup>de Naurois (1969a). <sup>4</sup>Supposedly reported by early authors fide Bourne (1955) but documentation unclear. <sup>5</sup>Interpreted here as this species, because reported to take poultry and catch fish at the coast (Keulemans, 1866), although interpreted by Hazevoet (1995) as red kite. <sup>6</sup>Keulemans (1866). <sup>7</sup>de Naurois (1987a). <sup>8</sup>Alexander (1898b). <sup>9</sup>Carter (2001). <sup>10</sup>van der Molen (2008). <sup>11</sup>Hazevoet (1995). <sup>12</sup>SMH personal records. <sup>13</sup>Breider et al. (2004). <sup>14</sup>No indication of numbers but all assumed to be visitors only. <sup>15</sup>Bannerman (1968). <sup>16</sup>Hazevoet (2003). <sup>17</sup>Bourne (1955). <sup>18</sup>Hille & Collar (2009). <sup>19</sup>Summers-Smith (1984). <sup>20</sup>Presumed vagrant, 1897 (Hazevoet, 1995). <sup>21</sup>Hille (1998). <sup>22</sup>Hille & Thiollay (2000). <sup>23</sup>Bolle (1856). <sup>24</sup>Salvadori (1899). <sup>25</sup>Apparently with a 50% decline. <sup>26</sup>Anderson & White (2000). <sup>27</sup>de Naurois (1973). <sup>28</sup>Hazevoet (1997). <sup>29</sup>Attribution of genetic material to bird from Boavista (Kruckenhauser et al., 2004) is a mistake for Santo Antão (SMH). <sup>30</sup>Krabbe et al. (2003). <sup>31</sup>Palma et al. (2004). <sup>32</sup>Ferreira & Palma (2000). <sup>33</sup>de Naurois (1987b). <sup>34</sup>Apparently 50% increase. <sup>35</sup>de Naurois & Bonnaffoux (1969). <sup>36</sup>Lopes Suarez & Varo Cruz (2007). <sup>37</sup>Apparently 40% decline. <sup>38</sup>Estimate of 77 (Ontiveros, 2005) was derived from inappropriate methodology and analysis. <sup>39</sup>de Naurois (1969b). <sup>40</sup>den Hartog (1990).

## Results

Scavenging raptors in Cape Verde have declined more steeply and across more islands than predatory raptors, with the partial scavenger, common buzzard, showing an intermediate decline between the mainly scavenging and the purely predatory species (Fig. 2; Table 1). Degree of change in status varies between islands, with Santo Antão offering the best prospects for conservation (Fig. 3). The literature and our data provide further more generalized evidence on the status of individual species.

*Black and red kites* Early visitors did not realize that both species, which are very similar, were present, so the record up to the 1950s is confused, with black kite being particularly poorly documented (Hille & Collar, 2009). Despite 'apparently identical ecologies' (Naurois, 1969a), the two may have diverged by habitat, with red kite more in mountains and black kite more on coasts (Hartog, 1990), which might explain why *fasciicauda* perhaps never occurred on low-lying Sal, Boavista and Maio (Hazevoet, 1995). However, the future of both species in Cape Verde is in doubt, with the black kite suffering a greater spread of severe declines than any other raptor (Fig. 2) and the red kite extinct or near-extinct (Table 1; Hille & Collar, 2009). The last red kites seen on Santo Antão, in the 1990s, were chasing young chickens in the mountains, hunting big grasshoppers and patrolling harbours, seeking food even




FIG. 2 Trends in populations of the seven raptor species across 13 Cape Verde islands (Fig. 1). Each island is assigned a trend category (severe decline, decline, or no change) for each species, and columns show the proportion of each trend category per species. No change indicates populations with R-R and C-C but not X?-X? (abbreviations defined in Table 1) because no viable population was recorded. C-R, R-X and R-X? (one-category changes) indicate decline, and C-X and C-X? (two-category changes) indicate severe decline.

near houses but always at a distance from people (SMH, pers. obs.). The population of the red kite may never have numbered > 100. In the late 1980s it was guessed to be 50–100 birds (including hybrids) and in the mid-1990s 'some tens of pairs' (Ferguson-Lees & Christie, 2001) but it is no longer certain that any survive (Hille & Collar, 2009).

*Egyptian vulture* This species ranges in dry open country from shoreline to high mountains (Hazevoet, 1995; SMH, pers. obs.). Cape Verde was not colonized by humans and their livestock until 1460 and never had an indigenous non-volant mammal, so any pre-settlement vulture must have subsisted on seabird and turtle colonies (Hartog, 1990); Naurois (1969a) noted that the species could catch grounded petrels near their burrows. Its scavenging habits earned it both legal protection (Bolle, 1856) and popular support (Bannerman, 1968) and in the early 1950s it was 'abundant around the large towns' except on Fogo and Brava, with 'one or two pairs around most villages' (Bourne, 1955). In the 1960s Naurois (1969a) reported that it 'swarms' (*pullule*) in the archipelago. However, by 1986–1993, never more than five were seen together anywhere except on Santo Antão (Hazevoet, 1995) and the species was 'declining at an alarming rate' (Hazevoet et al., 1996).

*Common buzzard* Largely found in remote montane areas, this typically soaring raptor, which takes both live and dead prey, went unrecorded by most pre-1950 explorers, only one of whom referred to it as 'not uncommon' (Bannerman, 1968). Hazevoet (1995, 1998, 2003) described it as 'exceedingly rare in last few decades', with a total population of 'some tens of pairs', and our analysis suggests declines on over half the islands. Records from Boavista comprise two




FIG. 3 Population trend and number of raptor species by island in Cape Verde. The index of change sums all changes for all species per island, obtained by subtracting 1 for one step down and 2 for two steps down between the categories severe decline, decline and no change (explained in Fig. 2). Santo Antão has the most species but shows the second slowest rate of deterioration, and is thus a conservation priority.

specimens and one recent sighting: James (1984) attributed the first specimen to long-legged buzzard *B. rufinus cirtensis*, and Hazevoet (1995, 2003) considered the second very similar to the first, so it is conceivable that the tiny population on this easternmost island actually involves a different species.

*Osprey* This all-year resident feeds exclusively on marine fish and nests from beaches to high mountains (SMH, pers. obs.). Bolle (1856) recorded it on 'all 10 islands' and as 'especially numerous on the eastern flatter sides where the sea is richest in fish' (our translation). Keulemans (1866) also described it as numerous. However, Alexander (1898b: 100) and the collectors on the *Blossom* (early 1920s) found it absent or very rare in the southern islands (Bourne, 1955); a circumstance confirmed by SMH and Palma et al. (2004). Naurois (1969a) estimated c. 50 pairs in the 1960s, which he considered reflected a decline attributable to human nest despoliation. Hazevoet (1995) also estimated c. 50 pairs in 1988–1993, with 3–8 pairs on each island, 1–2 pairs on each islet and largest numbers on Santo Antão and the three eastern islands. Surveys in 1996–1997 (SMH, pers. obs.) and 1998–2001 (Ferreira & Palma, 2000; Palma et al., 2004) resulted in estimates of pairs of 55–65 (SMH, pers. obs.), 44 or perhaps 60 (Ferreira & Palma, 2000), 76–86 (Barone et al., 2000) and 72–81 (Palma et al., 2004). Accessible sites in the 1960s and earlier have been abandoned and the species only survives in remoter areas (SMH, pers. obs.; Ferreira & Palma, 2000).

*Common kestrel* Two endemic subspecies inhabit rocky areas with slopes and updraughts. On the southern islands in 1897 'numerous pairs [could] frequently be found in one [palm] grove' (Alexander, 1898a), something not observed today (SMH, pers. obs.). Bannerman (1968) believed numbers were 'declining at a rapid rate' in the 1960s but intensive surveys on all islands in 1996–2001 (Hille et al., 2003) suggested no significant status change, yielding totals of 706 pairs of *neglectus* and 951 pairs of *alexandri* (SMH, pers. obs.; Table 1).

*Peregrine* Breeding was only proved in 1963 for this inhabitant of high cliffs and mountains on coasts and in island interiors (Bannerman, 1968; Naurois, 1969b). Naurois (1969a) thought it might be almost extinct and later argued, dubiously, that numbers must have been higher 60 years earlier, given the ease with which the *Blossom* expedition obtained specimens (Naurois, 1984). Hazevoet (1995) thought it probably occurred 'in very small numbers on all islands and islets', perhaps < 20 birds, later suggesting 'a few tens of pairs' (Hazevoet 1997, 2003). There has been virtually no status change.

## Discussion

The evidence from Cape Verde indicates that scavenging is a strategy in which the balance of risk and advantage is

rapidly lost under the influence of modernized societies (Fig. 2). Black and red kite populations have declined on all islands to the point of near-extinction. The Egyptian vulture has also suffered widespread declines, with only two populations, on Boavista and Santo Antão, categorized as no change (i.e. remaining in the category common), although the former is tiny and the latter has declined by 50%. The part-scavenging common buzzard showed a decrease on > 70% of islands, and only Santo Antão and Santiago harbour small but stable (no change) populations, although again the latter declined by > 50%. Ospreys also exhibited declines but continue to flourish in the north-west. Only common kestrel (both subspecies, all populations) has not changed in status, and nearly all peregrine populations remained stable.

Raptor extinctions on Mediterranean and Macaronesian islands have been shown to be more likely where human populations are denser, probably because of persecution, and species dependent on human activities are also more liable to extinction (Donazar et al., 2005). In Cape Verde the human population density increased from 36 km<sup>-2</sup> in 1950 to 109 km<sup>-2</sup> in 2000 (UN Department of Economic and Social Affairs, 2008). We outline the three major threats to raptors on the islands, in their likely order of impact.

*Pesticides and poisons* Hazevoet et al. (1996) and Hazevoet (1998, 1999) believed the decline of the red kite and common buzzard to be a side effect of the relatively modern local practice of poisoning feral dogs with laced meat, the birds eating either the bait or the corpses and, if not succumbing directly, then probably suffering decreased fertility. This problem also afflicts the Egyptian vulture, two bodies of which were found beside a dead cat at a Santo Antão dump in 2000 (SMH, pers. obs.). Rodenticides and insecticides may, however, also have significant effects. Pesticide use on the islands is common and includes organophosphorous substances. Farmers use carbamates as insecticides, nematocides, DDT, strychnine and rodenticides (e.g. Raticide, Racumin and Fostrien; SMH, pers. obs.). Fostrien was widely used on Santiago at least until the 1990s, and methyl parathion (E 601), which poses a higher risk to birds than to other vertebrates (Grue et al., 1983; Mañosa et al., 2003), has been used against grasshoppers in crop fields on some islands, including Boavista (SMH, pers. obs.).

*Persecution* Although Cape Verde children tend to persecute all raptors (SMH, pers. obs.), the two falcons and vulture generally escape human interference: the kestrel is harmless, the peregrine reclusive and the vulture useful (which perhaps helps explain why the vulture has fared slightly better than the other scavengers; Fig. 2). However, the two kites and buzzard have always attracted hostility as chicken thieves (Alexander, 1898b; Bannerman, 1968); even in the 1990s black kites would be pursued with stones

(SMH, pers. obs.). Kite nests were plundered, explaining the black kite's near-disappearance from São Vicente (Bannerman, 1968). Keulemans (1866) reported that local people regarded kite meat (especially nestlings) as a delicacy. The tradition has persisted, although interviewees today report disliking the flavour (Hille, 1998; SMH, pers. obs.). Eggs and young of ospreys were also considered delicacies: Naurois (1969a, 1984) blamed the species' decline on this, as did Ferreira & Palma (2000) for the decline on Sal, and thought the birds only escaped by nesting on inaccessible cliffs.

**Food supply decrease** Natural fluctuations in food supplies, caused by periodic drought, are common in Cape Verde, and predator populations fluctuate in response. Darwin's failure to record a single Egyptian vulture while prospecting Santiago in 1832, and breeding inactivity in common buzzards on Santiago in 1969, were attributed to drought-induced loss of, respectively, livestock (Dohrn, 1871; Bannerman, 1968) and rodents (Naurois, 1969a). However, in the past 150 years overall food availability has probably declined. Lower rainfall caused by Saharan expansion (Rosenfeld et al., 2001) and overgrazing by, and harvesting wild plants as fodder for, goats, which increased in numbers from 40,000 in 1961 to 148,000 in 2004 (República de Cabo Verde, 2004), must have produced much sparser, less productive vegetation. Invertebrate density has been thinned in farmed areas. Changes in livestock management (fewer available corpses and afterbirths), food resourcing (beef imported rather than reared locally), slaughterhouse practices (less waste), refuse disposal patterns (use of burning) and sanitation regimes (cleaner streets and flush toilets) have resulted in fewer carcasses, rodents and faeces (Bannerman, 1968; Hazevoet, 1995). There is no direct evidence of the impact of these various constraints but all five black kites caught on Maio in 2002 (Hille & Collar, 2009) showed stress bars in tail and wing feathers (SMH, pers. obs.).

Santo Antão, with its high agricultural output, permanent streams and comparatively low human density, is the stronghold of all raptor taxa in Cape Verde except the kites (Fig. 3). Its northern valleys, notably Ribeira da Torre, Ribeira Grande and Ribeira do Paul, harbour the last significant buzzard and vulture populations, and the highest densities of kestrels, in Cape Verde, and ospreys nest there close to the sea. Cape Verde is an Endemic Bird Area (Stattersfield et al., 1998) but although 12 Important Bird Areas have been identified there (Hazevoet, 2001), none is on Santo Antão. A protected area, with so-called restaurants to provide safe food for scavengers, as advocated in Spain by Hernández & Margalida (2009), coupled with a public awareness campaign, would be a valuable conservation measure in the island's northern valleys, especially in light of the vulture and buzzard declines on the island, which

indicate that land-use change and agricultural and other human practices continue to have a negative impact.

The plight of scavenging raptors in Cape Verde shows that this guild is significantly disadvantaged by human behaviour and economic development. It also points to the general conservation problem of scavengers on islands elsewhere in Macaronesia and the Mediterranean as a result of their naturally small populations and low rates of dispersal (Eurasian griffons, for example, have great difficulty crossing water; Bildstein et al., 2009). There is, moreover, emerging evidence that these populations have been isolated long enough to have developed into distinguishable demes and taxonomic entities. The Egyptian vultures on the Canaries are morphologically, genetically and hence taxonomically distinct (Donazar et al., 2002; Kretzmann et al., 2003), and even the seemingly undifferentiated black kite of Cape Verde possesses certain minor but clear morphological characters (Hille & Collar, 2009). We infer from these pieces of evidence that insular populations of scavenging raptors may represent locally adapted forms that cannot easily be replaced and therefore require targeted intervention to ensure their survival. Greater control of toxic materials, broad public awareness campaigns and the use of restaurants, which are also relevant to many continental populations, are crucial in this regard.

In October 2010 the European Union, after stalling for some years owing to public health considerations relating to TSE (transmissible spongiform encephalopathies), gave its approval to the use of feeding stations for scavenging raptors in all relevant range states (EC regulation 1069/2009). This welcome news unfortunately does not extend to Cape Verde, and the next step there will be to negotiate for a major landscape management initiative that will stabilize and improve the status of the country's most vulnerable raptors.

## Acknowledgements

SMH's fieldwork was funded by the German Ornithological Society, European Commission (Marie Curie grant) and Peregrine Fund. We thank Roy Dennis, Simon Thomsett, Jim Wilmwarth, Sandra Krätschmer, Ute Nau, Armin Nemitz and Gernot Segelbacher for help in the field, Maria Tereza Vera Cruz and Hans-Uwe Heckel for logistic support, K.L. Bildstein for helpful information, and J.A. Donazar and V. Bretagnolle for valuable comments.

## References

- ALEXANDER, B. (1898a) Further notes on the ornithology of the Cape Verde islands. *Ibis*, 4, 277–285.
- ALEXANDER, B. (1898b) An ornithological expedition to the Cape Verde islands. *Ibis*, 4, 74–118.

- ANDERSON, C.M. & WHITE, C.M. (2000) Recent observations on peregrine falcons *Falco peregrinus* of the Cape Verde islands, Atlantic Ocean. In *Raptors at Risk: Proceedings of the Fifth World Conference on Birds of Prey and Owls, Johannesburg, South Africa* (eds R.D. Chancellor & B.-U. Meyburg), pp. 685–689. Hancock House Publishers, Surrey, British Columbia, Canada.
- BANNERMAN, D.A. (1968) Bird species accounts. In *Birds of the Atlantic Islands* (eds D.A. Bannerman & W.M. Bannerman), p. 2. Oliver & Boyd, Edinburgh & London, UK.
- BARONE, T.R., DELGADO, C.G. & FERNÁNDEZ DEL CASTILLO, A.M. (2000) La avifauna nidificante del archipiélago de Cabo Verde. *Makaronesia, Boletín de la Asociación de Amigos del Museo de Ciencias Naturales de Tenerife*, 2, 43–55.
- BILDSTEIN, K.L., BECHARD, M.J., FARMER, C. & NEWCOMB, L. (2009) Narrow sea crossings present major obstacles to migrating griffon vultures *Gyps fulvus*. *Ibis*, 151, 382–391.
- BIRDLIFE INTERNATIONAL (2009) Species factsheet: *Milvus milvus*. <http://www.birdlife.org> [accessed 22 September 2009].
- BOLLE, C. (1856) Die Vogelwelt auf den Inseln des grünen Vorgebirges. *Journal of Ornithology*, 4, 17–31.
- BOURNE, W.R.P. (1955) The birds of the Cape Verde islands. *Ibis*, 97, 508–556.
- BREIDER, J.-B., FORSBERG, P., SVENSSON, A., ULLMAN, M. & WIGERMO, C. (2004) *Kap Verde Bird Report*. [http://www.club300.se/Files/TravelReports/KapVerde2004\\_JMB.pdf](http://www.club300.se/Files/TravelReports/KapVerde2004_JMB.pdf) [accessed 20 December 2009].
- CARTER, P. (2001) *Birding Trip Report: Cape Verde Islands, 1st Nov. to 5th Dec. 2001*. Unpublished Report.
- CUTHBERT, R., GREEN, R.E., RANADE, S., SARAVANAN, S., PAIN, D.J., PRAKASH, V. & CUNNINGHAM, A.V. (2006) Rapid population declines of Egyptian vulture (*Neophron percnopterus*) and red-headed vulture (*Sarcogyps calvus*) in India. *Animal Conservation*, 9, 349–354.
- DOHRN, H. (1871) Beiträge zur Ornithologie der Capverdischen Inseln. *Journal of Ornithology*, 19, 1–10.
- DONÁZAR, J.A., GANGOSO, L., FORERO, M.G. & JUSTE, J. (2005) Presence, richness and extinction of birds of prey in the Mediterranean and Macaronesian islands. *Journal of Biogeography*, 32, 1701–1713.
- DONÁZAR, J.A., NEGRO, J.J., PALACIOS, C.J., GANGOSO, L., GODOY, J.A., CEBALLOS, O. et al. (2002) Description of a new subspecies of the Egyptian vulture (Accipitridae: *Neophron percnopterus*) from the Canary Islands. *Journal of Raptor Research*, 36, 17–23.
- FERGUSON-LEES, I.J. & CHRISTIE, D.A. (2001) *Raptors of the World*. Christopher Helm, London, UK.
- FERREIRA, J. & PALMA, L. (2000) The osprey *Pandion haliaetus* in the Cape Verde islands: distribution, population trends and conservation problems. In *Raptors at Risk: Proceedings of the Fifth World Conference on Birds of Prey and Owls* (eds R.D. Chancellor & B.-U. Meyburg), pp. 721–727. Hancock House Publishers, Surrey, British Columbia, Canada.
- GRUE, C.E., FLEMING, W.J., BUSBY, D.G. & HILL, E.F. (1983) Assessing hazards of organophosphate pesticides to wildlife. *Transactions of the North American Wildlife and Natural Resources Conference*, 48, 200–220.
- HARTOG, J.C. DEN (1990) Birds of the Cape Verde islands. Notes on species observed (9 August–10 September 1986): distribution, migration, status, origin and conservation. In *8. Beitrag zur Fauna und Flora der Kapverdischen Inseln: Ergebnisse des 4. Symposiums* (ed. W. Lobin), pp.159–190. Senckenbergische Naturforschende Gesellschaft, Frankfurt-am-Main, Germany.
- HAZEVOET, C.J. (1995) *The Birds of the Cape Verde Islands*. British Ornithologists' Union (Check-list 13), Tring, UK.
- HAZEVOET, C.J. (1997) Notes on distribution, conservation, and taxonomy of birds from the Cape Verde islands, including records of six species new to the archipelago. *Bulletin of the Zoological Museum of the University of Amsterdam*, 15, 89–100.
- HAZEVOET, C.J. (1998) Third annual report on birds from the Cape Verde islands, including records of seven taxa new to the archipelago. *Bulletin of the Zoological Museum of the University of Amsterdam*, 16, 65–71.
- HAZEVOET, C.J. (1999) Notes on birds from the Cape Verde islands in the collection of the Centro de Zoologia, Lisbon, with comments on taxonomy and distribution. *Bulletin of the British Ornithologists' Club*, 119, 25–31.
- HAZEVOET, C.J. (2001) Cape Verde. In *Important Bird Areas in Africa and Associated Islands: Priority Sites for Conservation* (eds L.D.C. Fishpool & M.I. Evans), pp.161–168. Pisces Publications and BirdLife International (Conservation Series 11), Newbury and Cambridge, UK.
- HAZEVOET, C.J. (2003) Fifth report on birds from the Cape Verde islands, including records of 15 new taxa to the archipelago. *Arquivos do Museu Bocage*, NS, 3, 503–528.
- HAZEVOET, C.J., FISCHER, S. & DELOISON, G. (1996) Ornithological news from the Cape Verde islands in 1995, including records of species new to the archipelago. *Bulletin of the Zoological Museum of the University of Amsterdam*, 15, 21–27.
- HERNÁNDEZ, M. & MARGALIDA, A. (2009) Poison-related mortality effects in the endangered Egyptian vulture (*Neophron percnopterus*) population in Spain. *European Journal of Wildlife Research*, 55, 415–423.
- HILLE, S. (1998) Zur Situation der Milane *Milvus milvus fasciicauda* (Hartert, 1914) und *Milvus m. migrans* (Boddaert, 1783) auf den Kapverdischen Inseln. *Journal of Ornithology*, 139, 73–75.
- HILLE, S.M. & COLLAR, N.J. (2009) The taxonomic and conservation status of *Milvus* kites in the Cape Verde archipelago: further (and final?) reflections. *Bulletin of the British Ornithologists' Club*, 129, 217–221.
- HILLE, S.M., NESJE, M. & SEGELBACHER, G. (2003) Genetic structure of kestrel populations and colonization of the Cape Verde archipelago. *Molecular Ecology*, 12, 2145–2151.
- HILLE, S. & THIOLLAY, J.-M. (2000) The imminent extinction of the kites *Milvus milvus fasciicauda* and *Milvus m. migrans* on the Cape Verde islands. *Bird Conservation International*, 10, 361–369.
- JAMES, A.H. (1984) Geographic variation in the buzzard *Buteo buteo* (Linnaeus, 1758): mid-Atlantic and west Mediterranean islands (Aves: Accipitridae). *Beaufortia*, 34, 101–115.
- KEULEMANS, J.G. (1866) Opmerkingen over de vogels van de Kaap-Verdische Eilanden en van Prins-Eiland. *Nederlandsch Tijdschrift voor de Dierkunde*, 3, 363–374.
- KRABBE, E., ELIAS, G. & RILEY, A. (2003) *Bird Observations in the Cape Verde islands, 18–26 October 2003*. Unpublished Report.
- KRETZMANN, M.B., CAPOTE, N., GAUTSCHI, B., GODOY, J.A., DONÁZAR, J.A. & NEGRO, J.J. (2003) Genetically distinct island populations of the Egyptian vulture (*Neophron percnopterus*). *Conservation Genetics*, 4, 697–706.
- KRUCKENHAUSER, L., HARING, E., PINSKER, W., RIESING, M.J., WINKLER, H., WINK, M. & GAMAUF, A. (2004) Genetic vs morphological differentiation of Old World buzzards (genus *Buteo*, Accipitridae). *Zoologica Scripta*, 33, 197–211.
- LOPES SUAREZ, P. & VARO CRUZ, N. (2007) Preliminary data on distribution, population size, breeding activity and conservation problems of ospreys at Boa Vista (Cape Verde islands). *International Meeting on Western Palearctic Osprey Populations, 1–2 December 2007*. Alberese, Grosseto, Italy.
- MAÑOSA, S., MATEO, R., FREIXA, C. & GUITART, R. (2003) Persistent organochlorine contaminants in eggs of northern

- goshawk and Eurasian buzzard from north-eastern Spain: temporal trends related to changes in the diet. *Environmental Pollution*, 122, 351–359.
- MARTÍN, A. & LORENZO, J.A. (2001) *Aves del archipiélago canario*. Francisco Lemus, La Laguna, Tenerife, Spain.
- MOLEN, D. VAN DER (2008) *Cape Verde Islands, 21 February–2 March [2008]*. <http://www.birdtours.co.uk/tripreports/cape-verde> [accessed 28 September 2009].
- NAUROIS, R. DE (1969a) Notes brèves sur l'avifaune de l'archipel du Cap-Vert. Faunistique, endémisme, écologie. *Bulletin de l'Institut fondamental d'Afrique noire*, 31, 143–218.
- NAUROIS, R. DE (1969b) La population des Faucons pèlerins (*Falco peregrinus madens* Ripley et Watson) de l'archipel du Cap Vert: effectif, écologie et signification zoogéographique. *Alauda*, 37, 301–314.
- NAUROIS, R. DE (1973) Recherches sur la Buse (*Buteo buteo* L.) de l'archipel du Cap Vert. In *Livro de Homenagem ao Prof. Fernando Frade*, pp. 157–175. Junta de Investigações do Ultramar, Lisbon, Portugal.
- NAUROIS, R. DE (1984) Contribution à l'ornithologie de l'archipel du Cap Vert: reproducteurs menacés d'extinction, nicheurs occasionnels, oiseaux mentionnés par erreur. *Boletim do Museu Municipal do Funchal*, 36, 38–50.
- NAUROIS, R. DE (1987a) Les oiseaux de l'archipel du Cap Vert. *Bulletin de la Société Zoologique de France*, 112, 307–326.
- NAUROIS, R. DE (1987b) Le Balbuzard (*Pandion haliaetus* L.) aux îles du Cap Vert. *Annali del Museo Civico di Storia Naturale di Genova*, 86, 657–681.
- NAUROIS, R. DE & BONNAFFOUX, D. (1969) L'avifaune de l'île du Sel (Ilha do Sal, Archipel du Cap Vert). *Alauda*, 37, 94–113.
- NØRREVANG, A. & HARTOG, J.C. DEN (1984) Bird observations in the Cape Verde islands (4–22 June 1982). In *Ergebnisse des 2. Symposiums 'Fauna und Flora der Kapverdischen Inseln'* (ed. W. Lobin), pp. 107–134. Senckenbergische Naturforschende Gesellschaft, Frankfurt am Main, Germany.
- ONTIVEROS, D. (2005) Abundance and diet of Alexander's kestrel (*Falco tinnunculus alexandri*) on Boavista island (Archipelago of Cape Verde). *Journal of Raptor Research*, 39, 82–85.
- PAIN, D.J., BOWDEN, C.G.R., CUNNINGHAM, A.V., CUTHBERT, R., DAS, D., GILBERT, M. et al. (2008) The race to prevent the extinction of South Asian vultures. *Bird Conservation International*, 18, S30–S48.
- PALMA, L., FERREIRA, J., CANGARATO, R. & VAZ PINTO, P. (2004) Current status of the osprey in the Cape Verde islands. *Journal of Raptor Research*, 38, 141–147.
- REPÚBLICA DE CABO VERDE (2004) *Recenseamento geral da agricultura 2004—resultados definitivos*. Ministério do Ambiente, Agricultura e Pescas. [http://www.fao.org/fileadmin/templates/ess/documents/world\\_census\\_of\\_agriculture/main\\_results\\_by\\_country/Capeverde\\_2004F\\_1.pdf](http://www.fao.org/fileadmin/templates/ess/documents/world_census_of_agriculture/main_results_by_country/Capeverde_2004F_1.pdf) [accessed 3 October 2009].
- ROSENFELD, D., RUDICH, Y. & LAHAV, R. (2001) Desert dust suppressing precipitation—a possible desertification feedback loop. *Proceedings of the National Academy of Sciences*, 98, 5975–5980.
- SALVADORI, T. (1899) Collezioni ornitologiche fatte nelle Isole del Capo Verde da Leonardo Fea. *Annali del Museo Civico di Storia Naturale di Genova*, 20, 283–312.
- SEKERCIOGLU, C.H., DAILY, G.C. & EHRLICH, P.R. (2004) Ecosystem consequences of bird declines. *Proceedings of the National Academy of Sciences*, 101, 18042–18047.
- SERGIO, F., PEDRINI, P. & MARCHESI, L. (2003) Reconciling the dichotomy between single species and ecosystem conservation: black kites (*Milvus migrans*) and eutrophication in pre-Alpine lakes. *Biological Conservation*, 110, 101–111.
- STATTERSFIELD, A.J., CROSBY, M.J., LONG, A.J. & WEGE, D.C. (1998) *Endemic Bird Areas of the World: Priorities for Biodiversity Conservation*. BirdLife International (Conservation Series 7), Cambridge, UK.
- SUMMERS-SMITH, D. (1984) Bird notes from the Cape Verde islands. *Bulletin of the British Ornithologists' Club*, 104, 48–149.
- THIOLLAY, J.-M. (2006) The decline of raptors in West Africa: long-term assessment and the role of protected areas. *Ibis*, 148, 240–254.
- TUCKER, G.M. & HEATH, M.F. (1994) *Birds in Europe: Their Conservation Status*. BirdLife International (Conservation Series 3), Cambridge, UK.
- UN DEPARTMENT OF ECONOMIC AND SOCIAL AFFAIRS (2008) *World Population Prospects: The 2008 Revision*. <http://esa.un.org/unpp> [accessed 2 November 2010].
- XIROUCHAKIS, S. & TSIKIRIS, R. (2009) Situación y tendencias poblacionales de los buitres en Grecia. *Munibe*, 29, 155–171.

## Biographical sketches

SABINE M. HILLE studies the evolution and dynamics of small bird populations on islands and in changing environments. She has been doing research in Cape Verde for several years. NIGEL J. COLLAR works on threatened bird species, running projects in Angola and Ethiopia.