

ICONES PLANTARUM;

OR

FIGURES,

WITH

BRIEF DESCRIPTIVE CHARACTERS AND REMARKS,

OF

NEW OR RARE PLANTS,

SELECTED FROM THE AUTHOR'S HERBARIUM.

By SIR WILLIAM JACKSON HOOKER, K.H.,

LL.D., F.R.S., AND F.L.S.

VICE-PRESIDENT OF THE LINNEAN SOCIETY,

MEMBER OF THE IMP. ACAD. NAT. CUR., ETC., ETC., ETC.

HONORARY MEMBER OF THE ROYAL IRISH ACADEMY, OF THE ROYAL MEDICAL AND
SURGICAL SOC. OF LONDON, ETC., ETC.

AND

DIRECTOR OF THE ROYAL BOTANICAL GARDENS, SER.

VOL. IV. NEW SERIES,

OR VOL. VIII. OF THE ENTIRE WORK.

LONDON:

HIPPOLYTE BAILLIÈRE,

FOREIGN BOOKSELLER TO THE ROYAL COLLEGE OF SURGEONS, AND TO THE ROYAL
MEDICAL AND SURGICAL SOCIETY.

219, REGENT STREET.

PARIS: J. B. BAILLIÈRE, RUE DE L'ÉCOLE DE MÉDECINE.

1848.

TAB. DCCLIV.

LOTUS BRUNNERI, *Webb*.

Caulē fruticoso procumbente, foliis pube cinerea appressa sericeis, stipulis lanceolatis sessilibus petiolo brevioribus caducis, foliolis obovato-lanceolatis, pedunculis paucifloris, calycibus urceolato-campanulatis sericeo-pubescentibus, dentibus lanceolatis acutis, corolla lutea, vexillo elliptico subacuto carina sublongiore. *Webb*.

Lotus Brunneri. *Webb, Spicil. Gorgon. in Hook. Niger Voy.* p. 119, Tab. III.

Lotus anthylloides? *Brunn. Ergebn. p. 86, (non Vent.)*

HAB. Island of Sal, Cape de Verd. *Brunner*.

A plant perfectly distinct from the *L. anthylloides*, Vent. It differs from *L. Jacobæus* by the form of its leaves, by its appressed, and not villous and patent, pubescence, by its short sessile stipules, by the teeth of the calyx being merely acute, and not more or less filiform at the extremity, and by the form and colour of the corolla. *Webb*.

Fig. 1. Flower. *f. 2.* Vexillum. *f. 3.* Ala. *f. 4.* Carina.
f. 5. Stamina and ovarium. *f. 6.* Ovarium. *f. 7.* Pod.
f. 8. Seed:—magnified.

